

NDC 51660-351-05
Maximum Strength

Ranitidine Tablets USP, 150 mg Acid Reducer

✔ Prevents & Relieves Heartburn Associated
 with Acid Indigestion & Sour Stomach

**THIS PACKAGE FOR HOUSEHOLDS
 WITHOUT YOUNG CHILDREN.**

500 TABLETS

IMPORTANT: Read the directions, consumer information leaflet and warnings before use. It contains important information.

Drug Facts

Active ingredient (in each tablet)
 Ranitidine 150 mg (as ranitidine hydrochloride USP, 168 mg).....Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Warnings

Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers
- if you have kidney disease, except under the advice and supervision of a doctor.

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions


- adults and children 12 years and over:
 - to relieve symptoms, swallow 1 tablet with a glass of water
 - to prevent symptoms, swallow 1 tablet with a glass of water **30 to 60 minutes before eating** food or drinking beverages that cause heartburn
 - can be used up to twice daily (do not take more than 2 tablets in 24 hours)
 - children under 12 years: ask a doctor

Other information

- store at 20° - 25° C (68° - 77° F)
- avoid excessive heat or humidity
- this product is sodium and sugar free
- **TAMPER EVIDENT: DO NOT USE IF THE PRINTED FOIL UNDER CAP IS OPEN OR TORN.**

Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, talc, titanium dioxide

Questions? call 1-800-406-7984



XXXXXX
 FPO
 0 000000000000 0

0415
 Distributed by:
 Ohm Laboratories Inc.
 1385 Livingston Avenue
 North Brunswick, NJ 08902
 MADE IN INDIA

Batch No. _____
 Expiration Date: _____

non varnish area

Standard Format
 Helvetica
 Variable horizontal scale
 adheres to 39 characters per inch

Drug Facts -10 pt. Helv. Bold Italic, cond 85%

Headings - 8 pt. Helv. Bld. Italic, cond 85%

Sub Heads - 6 pt. Helv. Bold, 85%


Text 6 pt. Helv., 85%, min. leading 5.54

Border 2.5 pt. rule, 1/8 gap all around

Square Bullets - 5 pt. Zapf Ding., cond 85%

Barline 2.5 pt. rule

CARTON



DESCRIPTION : **Ranitidine 150mg Tabs 500 Label**

DIMENSION : **8" x 3"**

SAP CODE : _____ OLD SAP CODE : _____

PROOF SIZE : **100%**

TRACKING : **AA 20/05/15, AA 22/05/15, AA26/05/15, AA27/05/15**

COLOR **BLACK**

Approved By : _____

Date : _____