

10"

Medication Guide available at:
www.strides.com/oxyco-hcl-tabs

Medication Guide
**Oxycodone Hydrochloride (ox” i koe’ done hye” droe
klor’ ide) Tablets USP, ©**

Oxycodone hydrochloride is:

- A strong prescription pain medicine that contains an opioid (narcotic) that is used to manage pain severe enough to require an opioid pain medicine, when other pain treatments such as non-opioid pain medicines do not treat your pain well enough or you cannot tolerate them.
- An opioid pain medicine that can put you at risk for overdose and death. Even if you take your dose correctly as prescribed you are at risk for opioid addiction, abuse, and misuse that can lead to death.

Important information about oxycodone hydrochloride:

- Get emergency help or call 911 right away if you take too much oxycodone hydrochloride (overdose).** When you first start taking oxycodone hydrochloride, when your dose is changed, or if you take too much (overdose), serious or life-threatening breathing problems that can lead to death may occur. Talk to your healthcare provider about naloxone, a medicine for the emergency treatment of an opioid overdose.
- Taking oxycodone hydrochloride with other opioid medicines, benzodiazepines, alcohol, or other central nervous system depressants (including street drugs) can cause severe drowsiness, decreased awareness, breathing problems, coma, and death.
- Never give anyone else your oxycodone hydrochloride. They could die from taking it. Selling or giving away oxycodone hydrochloride is against the law.
- Store oxycodone hydrochloride securely, out of sight and reach of children, and in a location not accessible by others, including visitors to the home.

Do not take oxycodone hydrochloride if you have:

- severe asthma, trouble breathing, or other lung problems.
- a bowel blockage or have narrowing of the stomach or intestines.
- allergy to oxycodone.

Before taking oxycodone hydrochloride, tell your healthcare provider if you have a history of:

- head injury, seizures
- problems urinating
- abuse of street or prescription drugs, alcohol addiction, opioid overdose, or mental health problems.
- liver, kidney, thyroid problems
- pancreas or gallbladder problems

Tell your healthcare provider if you are:

- noticing your pain getting worse.** If your pain gets worse after you take oxycodone hydrochloride, do not take more of oxycodone hydrochloride without first talking to your healthcare provider. Talk to your healthcare provider if the pain that you have increases, if you feel more sensitive to pain, or if you have new pain after taking oxycodone hydrochloride.
- pregnant or planning to become pregnant.** Use of oxycodone hydrochloride for an extended period of time during pregnancy can cause withdrawal symptoms in your newborn baby that could be life-threatening if not recognized and treated.
- breastfeeding.** Oxycodone hydrochloride passes into breast milk and may harm your baby. Carefully observe infants for increased sleepiness (more than usual), breathing difficulties, or limpness. Seek immediate medical care if you notice these signs.
- living in a household where there are small children or someone who has abused street or prescription drugs
- taking prescription or over-the-counter medicines, vitamins, or herbal supplements. Taking oxycodone hydrochloride with certain other medicines can cause serious side effects that could lead to death.

4"

Front Page

NON PRINTING COLOUR

MFG. LOCATION: CR Site

ARTWORK DETAIL LABEL

Product	Oxycodone HCL Tablets, USP				
Buyer/Country	STRIDES PHARMA INC - US		Component	Medication Guide	
Dimension	4"x 10"			Pack	--
New Item Code	1052534 (MC-028-01-02)		Old Item Code	NA	
Colour Shades	BLACK BLACK			No. of Colours	1
Change Control No.	CR-01238			Artwork Version	1.0
Design/Style	Front & Back Side Printing. To be supplied in the FOLDED BOOKLET FORM with pasting.				
Substrate	40 GSM Bible Paper.				
Special Instructions	PRINTING CLARITY TO BE CLEAR AND SHARP				
Autocartonator Requirements	NA				
Caution to the printer: Before processing, please ensure that the ARTWORK received for printing is exactly in line with APPROVED ARTWORK provided to you. In case of any FONTS/DESIGN are Mis-matching with the APPROVED ARTWORK, please inform PDC for further action. DO NOT MAKE ANY CHANGE TO THE ARTWORK WITHOUT WRITTEN INSTRUCTIONS FROM PDC.					

RD/PD/003/F-08/R0

10"

When taking oxycodone hydrochloride:

- Do not change your dose. Take oxycodone hydrochloride exactly as prescribed by your healthcare provider. Use the lowest dose possible for the shortest time needed.
- For acute (short-term) pain, you may only need to take oxycodone hydrochloride for a few days. You may have some oxycodone hydrochloride left over that you did not use. See disposal information at the bottom of this section for directions on how to safely throw away (dispose of) your unused oxycodone hydrochloride.
- Take your prescribed dose every 4 to 6 hours as needed for pain. Do not take more than your prescribed dose. If you miss a dose, take your next dose at your usual time.
- Call your healthcare provider if the dose you are taking does not control your pain.
- If you have been taking oxycodone hydrochloride regularly, do not stop taking oxycodone hydrochloride without talking to your healthcare provider.
- Dispose of expired, unwanted, or unused oxycodone hydrochloride by promptly flushing down the toilet, if a drug take-back option is not readily available. Visit www.fda.gov/drugdisposal for additional information on disposal of unused medicines.

While taking oxycodone hydrochloride DO NOT:

- Drive or operate heavy machinery, until you know how oxycodone hydrochloride affects you. Oxycodone hydrochloride can make you sleepy, dizzy, or lightheaded.
- Drink alcohol or use prescription or over-the-counter medicines that contain alcohol. Using products containing alcohol during treatment with oxycodone hydrochloride may cause you to overdose and die.

The possible side effects of oxycodone hydrochloride are:

- constipation, nausea, sleepiness, vomiting, tiredness, headache, dizziness, abdominal pain. Call your healthcare provider if you have any of these symptoms and they are severe.

Get emergency medical help or call 911 right away if you have:

- trouble breathing, shortness of breath, fast heartbeat, chest pain, swelling of your face, tongue, or throat, extreme drowsiness, light-headedness when changing positions, feeling faint, agitation, high body temperature, trouble walking, stiff muscles, or mental changes such as confusion.

These are not all the possible side effects of oxycodone hydrochloride. Call your healthcare provider for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. **For more information go to dailymed.nlm.nih.gov, www.strides.com or call 1-877-244-9825.**

This Medication Guide has been approved by the U.S. Food and Drug Administration.
Distributed by:
Strides Pharma Inc.
Bridgewater, NJ 08807

Revised: 01/2026
MC-028-01-02

Back Page

4"

NON PRINTING COLOUR

MFG. LOCATION: CR Site

ARTWORK DETAIL LABEL

Product	Oxycodone HCL Tablets, USP				
Buyer/Country	STRIDES PHARMA INC - US	Component	Medication Guide		
Dimension	4"x 10"			Pack	--
New Item Code	1052534 (MC-028-01-02)	Old Item Code	NA		
Colour Shades	 BLACK	No. of Colours			1
Change Control No.	CR-01238			Artwork Version	1.0
Design/Style	Front & Back Side Printing. To be supplied in the FOLDED BOOKLET FORM with pasting.				
Substrate	40 GSM Bible Paper.				
Special Instructions	PRINTING CLARITY TO BE CLEAR AND SHARP.				
Autocartonator Requirements	NA				
Caution to the printer: Before processing, please ensure that the ARTWORK received for printing is exactly in line with APPROVED ARTWORK provided to you. In case of any FONTS/DESIGN are Mis-matching with the APPROVED ARTWORK, please inform PDC for further action. DO NOT MAKE ANY CHANGE TO THE ARTWORK WITHOUT WRITTEN INSTRUCTIONS FROM PDC.					

RD/PD/003/F-08/R0