

(pol" ee eth' i leen glye' kol 3350, soe' dee um
 klor' ide, soe' dee um bye kar' bo nate and
 poe tas' ee um klor' ide)

What is the most important information I should know about polyethylene glycol 3350, sodium chloride, sodium bicarbonate and potassium chloride for oral solution?

Serious loss of body fluid (dehydration) and changes in blood salts (electrolytes) in your blood.

- abnormal heartbeats that may be a sign of a heart problem

- **seizures.** This can happen even if you have never had a seizure.
- **kidney problems**

- have heart problems
- have kidney problems
- take water pills or non-steroidal anti-inflammatory drugs (NSAIDs)

- vomiting that prevents you from keeping down the solution
- dizziness
- urinating less often than normal
- headache

What is polyethylene glycol 3350, sodium chloride, sodium bicarbonate and potassium chloride for oral solution?

Polyethylene glycol 3350, sodium chloride, sodium bicarbonate and potassium chloride for oral solution is safe and effective for use in pediatric patients aged 6 months and older.

Do not take polyethylene glycol 3350, sodium chloride, sodium bicarbonate and potassium chloride for oral

- a blockage in your bowel (obstruction)
- an opening in the wall of your stomach or intestine (bowel perforation)
- problems with food and fluid emptying from your stomach (gastric retention)
- a very dilated intestine (bowel)
- an allergy to any of the ingredients in polyethylene glycol 3350, sodium chloride, sodium bicarbonate and potassium chloride for oral solution. See the end of this leaflet for a complete list of ingredients in polyethylene glycol 3350, sodium chloride, sodium bicarbonate and potassium chloride for oral solution.

Before you take polyethylene glycol 3350, sodium chloride, sodium bicarbonate and potassium chloride for oral solution, tell your healthcare provider if you:

- have heart problems
- have stomach or bowel problems
- have ulcerative colitis
- have problems with swallowing or gastric reflux
- have a history of seizures
- are withdrawing from drinking alcohol
- have a low blood salt (sodium) level
- have kidney problems
- any other medical conditions
- are pregnant. It is not known if polyethylene glycol 3350, sodium chloride, sodium bicarbonate and potassium chloride for oral solution will harm your unborn baby. Talk to your doctor if you are pregnant or plan to become pregnant.
- are breastfeeding or plan to breastfeed. It is not known if polyethylene glycol 3350, sodium chloride, sodium bicarbonate and potassium chloride for oral solution passes into your breast milk. You and your healthcare provider should decide if you will take polyethylene glycol 3350, sodium chloride, sodium bicarbonate and potassium chloride for oral solution while breastfeeding.

Polyethylene glycol 3350, sodium chloride, sodium bicarbonate and potassium chloride for oral solution may affect how other medicines work. Medicines taken by mouth may not be absorbed properly when taken within 1 hour before the start of polyethylene glycol 3350, sodium chloride, sodium bicarbonate and potassium chloride for oral solution.

- medicines for blood pressure or heart problems
- medicines for kidney problems
- medicines for seizures
- water pills (diuretics)
- non-steroidal anti-inflammatory medicines (NSAID) pain medicines
- laxatives
- starch-based thickeners. For patients who have trouble swallowing, do not mix polyethylene glycol 3350, sodium chloride, sodium bicarbonate and potassium chloride for oral solution, with starch-based thickeners

Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist when you get a new medicine.

How should I take polyethylene glycol 3350, sodium chloride, sodium bicarbonate and potassium chloride for oral solution?

- Take polyethylene glycol 3350, sodium chloride, sodium bicarbonate and potassium chloride for oral solution exactly as your healthcare provider tells you to take it.
- Drink 240 mL (8 oz.) every 10 minutes. Rapid drinking of each portion is better than drinking small amounts.
- The first bowel movement should occur approximately one hour after you start drinking the solution.
- You may experience some abdominal bloating and distention before the bowels start to move. If severe discomfort or distention occur, stop drinking temporarily or drink each portion at longer intervals until the discomfort goes away.

- Continue drinking until the watery stool is clear and free of solid matter. This usually requires 3 liters and it is best to drink all of the solution.
- **Do not take undissolved polyethylene glycol 3350, sodium chloride, sodium bicarbonate and potassium chloride for oral solution's powder that has not been mixed with water (diluted), it may increase your risk of nausea, vomiting and fluid loss (dehydration).**
- Each jug of polyethylene glycol 3350, sodium chloride, sodium bicarbonate and potassium chloride for oral solution must be reconstituted with water (diluted) to 4 liters total volume before drinking.
- Do not take other laxatives while taking polyethylene glycol 3350, sodium chloride, sodium bicarbonate and potassium chloride for oral solution.
- **Do not eat solid foods on the day before your colonoscopy and until after your colonoscopy.** Drink only clear liquids:
 - the day before your colonoscopy
 - while taking polyethylene glycol 3350, sodium chloride, sodium bicarbonate and potassium chloride for oral solution
 - after taking polyethylene glycol 3350, sodium chloride, sodium bicarbonate and potassium chloride for oral solution until 2 hours before your colonoscopy

Polyethylene glycol 3350, sodium chloride, sodium bicarbonate and potassium chloride for oral solution can cause serious side effects, including:

- See Section “What is the most important information I should know about polyethylene glycol 3350, sodium chloride, sodium bicarbonate and potassium chloride for oral solution?”
- **changes in certain blood tests.** Your healthcare provider may do blood tests after you take polyethylene glycol 3350, sodium chloride, sodium bicarbonate and potassium chloride for oral solution to check your blood for changes. Tell your healthcare provider if you have any symptoms of too much fluid loss, including:
 - vomiting
 - nausea
 - bloating
 - dizziness
 - stomach (abdominal) cramping
 - headache
 - urinate less than usual
 - trouble drinking clear liquid
- **heart problems.** Polyethylene glycol 3350, sodium chloride, sodium bicarbonate and potassium chloride for oral solution may cause irregular heartbeats.
- **seizures**
- **ulcers of the bowel or bowel problems (ischemic**

The most common side effects of polyethylene glycol 3350, sodium chloride, sodium bicarbonate and potassium chloride for oral solution include:

- Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

Call your doctor for medical advice about side effects.
You may report side effects to FDA at 1-800-FDA-1088.

- Store polyethylene glycol 3350, sodium chloride, sodium bicarbonate and potassium chloride for oral solution at room temperature, between 59°F to 86°F (15°C to 30°C).

General information about the safe and effective use of polyethylene glycol 3350, sodium chloride, sodium bicarbonate and potassium chloride for oral solution.

This Medication Guide summarizes important information about polyethylene glycol 3350, sodium chloride, sodium bicarbonate and potassium chloride for oral solution. If you would like more information, talk with your healthcare provider. You can ask your pharmacist or healthcare provider for information that is written for healthcare professionals.

What are the ingredients in polyethylene glycol 3350, sodium chloride, sodium bicarbonate and potassium chloride for oral solution?

Inactive ingredients: Flavor lemon or flavor orange (in its respective flavor pack) with acesulfame potassium

Distributed by:
Strides Pharma Inc.
East Brunswick, NJ 08816

Revised: 10/2022

Revision: 6

Medication Guide available at:
www.strides.com/medication-guides

NOTE: Artwork is prepared for Electronic Medication Guide Purpose - Not to be printed.

ARTWORK DETAIL LABEL

Product	POLYETHYLENE GLYCOL 3350, SODIUM CHLORIDE, SODIUM BICARBONATE and POTASSIUM CHLORIDE for Oral Solution				
Buyer/Country	STRIDES PHARMA INC		Component	Medication Guide	
Dimension	200 x 335 mm			Pack	-
New Item Code	1047932		Old Item Code	1042926	
Colour Shades	Black			No. of Colours	1
Change Control No.	PC-TSG/2022/311 - Record Number: 359896			Artwork Version	7.0
Design/Style	NA				
Substrate	NA				
Special Instructions	NA				
Autocartanator Requirements	NA				
Caution to the printer: Before processing, please ensure that the ARTWORK received for printing is exactly in line with APPROVED ARTWORK provided to you. In case of any FONTS/DESIGN are Mis-matching with the APPROVED ARTWORK, please inform PDC for further action. DO NOT MAKE ANY CHANGE TO THE ARTWORK WITHOUT WRITTEN INSTRUCTIONS FROM PDC.					