

IBUPROFEN TABLETS USP 200 mg (Brown)

Final Carton Labeling for HDPE BOTTLE 50 Coated Caplets

(Package Child - Resistant)

Glue Area

See New Warnings Information NDC 57910-403-23

IBUPROFEN TABLETS USP

200 mg

Pain Reliever / Fever Reducer (NSAID)*

50 Coated Caplets**
**capsule-shaped tablets

*nonsteroidal anti-inflammatory drug

Glue Area

Open From This Side

Important: Read all product information before using.
Keep this box for important information.

LOT: _____
EXP: _____
Varnish free area

NDC 57910-403-23

Glue Area

Glue Area

Drug Facts (Continued)

- feel faint ■ vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.
In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**

adults and children 12 years and older	■ take 1 caplet every 4 to 6 hours while symptoms persist	■ if pain or fever does not respond to 1 caplet, 2 caplets may be used
	■ do not exceed 6 caplets in 24 hours, unless directed by a doctor	■ ask a doctor

Other Information

- Store between 20-25°C (68-77°F)

Drug Facts (Continued)

- tamper evident : do not use if imprinted safety seal under cap is broken or missing
- see end panel for lot number and expiry date

Inactive Ingredients

colloidal silicon dioxide, corn starch, hypromellose, magnesium stearate, microcrystalline cellulose, pregelatinized starch, red iron oxide, sodium starch glycolate, titanium dioxide, triacetin and talc

Questions or comments ?
Call : 1-732-465-0700
Weekdays 8.00 AM to 8.00 PM

Manufactured by:
Shasun Pharmaceuticals Limited
Unit-II, R.S.No: 32, 33, 34, Shasun Road
Periyakalpet, Pudukchery - 605 014,
INDIA Mfg.Lic.No. : 05 13 1523

Glue Area

Glue Area

Drug Facts (Continued)

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever / fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:

Glue Area

Artwork Scale: 44 mm x 44 mm x 75 mm			
Text Font type and size			
Font	: Arial		
Font Size			
Drug Facts	: 9 pt Bold Italic	Text	: 6 pt
Drug Facts (Continued)	: 8pt	Bullet	: 6 pt
Heading	: 7 pt Bold	Leading Space	: 6 pt
Sub Heading	: 6 pt Bold	Bar Line	: 1 pt
		Hair Line	: 0.5pt