

Glue Area

Drug Facts (continued)

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 - have 3 or more alcoholic drinks every day while using this product
 - take more or for a longer time than directed
- Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

- Do not use**
- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
 - right before or after heart surgery

- Ask a doctor before use if**
- you have problems or serious side effects from taking pain relievers or fever reducers
 - the stomach bleeding warning applies to you
 - you have a history of stomach problems, such as heartburn
 - you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
 - you are taking a diuretic

- Ask a doctor or pharmacist before use if you are**
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
 - under a doctor's care for any serious condition
 - taking any other drug

- When using this product**
- take with food or milk if stomach upset occurs

Glue Area

Important: Read all product information before using. Keep this box for important information.

Drug Facts

Active ingredient	Purpose
(in each orange caplet)	
Ibuprofen USP 200 mg (NSAID)* Pain reliever
*nonsteroidal anti-inflammatory drug	/fever reducer

- Uses**
- temporarily relieves minor aches and pains due to:
 - headache
 - minor pain of arthritis
 - backache
 - menstrual cramps
 - muscular aches
 - toothache
 - the common cold
 - temporarily reduces fever

- Warnings**
- Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.
- Symptoms may include:
- hives
 - facial swelling
 - shock
 - asthma (wheezing)
 - skin reddening
 - rash
 - blisters
- If an allergic reaction occurs, stop use and seek medical help right away.
- Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding.
- The chance is higher if you:
- are age 60 or older
 - have had stomach ulcers or bleeding problems
 - take a blood thinning (anticoagulant) or steroid drug

See New Warnings

NDC 59556-856-25

Drug Facts (continued)

Other information

- store between 20-25°C (68-77°F)
- do not use if imprinted text "SEALED for YOUR PROTECTION" on the safety seal under cap is broken or missing
- see end panel for lot number and expiration date
- Sodium free
- Each tablet contains 0.714 mg of Magnesium

Inactive ingredients

colloidal silicon dioxide, corn starch, hydroxypropylmethylcellulose, microcrystalline cellulose, pregelatinized starch, FD&C Yellow # 6 Al lake, sodium starch glycolate, titanium dioxide, Ineosin and talc

Questions or comments?

Call 1-855-742-7868 (toll-free)

Open From This Side

50 Coated Caplets**

**capsule-shaped tablets

*nonsteroidal anti-inflammatory drug

See New Warnings

NDC 59556-856-25

Ibuprofen Tablets, USP

200 mg

Pain Reliever/Fever Reducer (NSAID)*

Manufactured by:
Strides Shasun Limited
Puducherry - 605 014, India.
PON/DRUGS/16 13 4193

Distributed by:
Strides Pharma Inc.
East Brunswick, NJ 08816

Revised: 05/2017



3 15 9 5 5 6 18 5 6 2 5 1 9

Non Varnish Area

LOT :
EXP :

To be printed online

Drug Facts (continued)

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.(1-800-222-1222)

Directions

- do not take more than directed
- the smallest effective dose should be used

adults and children 12 years and older	<ul style="list-style-type: none"> ■ take 1 caplet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor
children under 12 years	<ul style="list-style-type: none"> ■ ask a doctor

Artwork Scale: 44 mm x 44 mm x 90 mm

Text Font type and size

Font	: Helvetica	
Font Size	: 9 pt Bold Italic	Text : 6 pt
Drug Facts	: 8pt	Bullet : 5 pt
Drug Facts (continued)	: 8 pt Bold	Leading Space : 7 pt
Heading	: 6 pt Bold	Bar Line : 2.5 pt
Sub Heading		Hair Line : 0.5pt